

Khon Kaen University

Gram negative bacilli in skin infection

PARAMA BUDMALA, MD.
CLINICAL PATHOLOGIST
DIVISION OF MICROBIOLOGY & IMMUNOLOGY
PARABUD@KKU.AC.TH



Khon Kaen University

Gram positive bacilli

PARAMA BUDMALA, MD.
CLINICAL PATHOLOGIST
DIVISION OF MICROBIOLOGY & IMMUNOLOGY
PARABUD@KKU.AC.TH



Bacterial skin infection

Objectives


1

- **Gram positive bacilli**

- *Bacillus anthracis*

- **Gram negative bacilli**


- *Pseudomonas aeruginosa*
- *Burkholderia pseudomallei*
- *Vibrio vulnificus*
- *Aeromonas hydrophila*



2

Concept for medical doctor

- Pathogen: Pathogenesis, virulence factor
- Clinical manifestation: history + physical examination
- Diagnosis criteria: clinical +/- laboratory
- Treatment and prevention



Bacillus spp.

Species

- *Bacillus cereus*: recooked rice and grain
- *Bacillus subtilis*: non-pathogenic in soil, water
- *Bacillus licheniformis*: Dairy product

Clinical significance

- Foodborne illness
 - Food poisoning

Diagnosis



- History taking

Treatment



- Supportive

+ Virulence factor

- Heat,acid-stable toxin (cereulide), emetic type
- Heat-labile Enterotoxin, diarrheal type eg. hemolysin BL (HBL), nonhemolytic enterotoxin (NHE), and cytotoxin K



Bacillus anthracis



Medusa head colony

Bacillus anthracis



A





B



C




D



Treatment

- Ciprofloxacin
 - Cutaneous: 7-10 days
 - Others: 60 days





A




B



C




D



Eschar

Cutaneous anthrax
DDx Rickettsial eschar



- Flu-like syndrome
- Hemoptysis, chest pain
- Mediastinitis/Hypoxia/ Dyspnea
- Lymphadenopathy

Pulmonary anthrax
(wool sorter's disease)



Gastrointestinal anthrax

Bacterial skin infection

Objectives


1

- **Gram positive bacilli**

- *Bacillus anthracis*

- **Gram negative bacilli**

- *Pseudomonas aeruginosa*
- *Burkholderia pseudomallei*
- *Vibrio vulnificus*
- *Aeromonas hydrophila*



2

Concept for medical doctor

- Pathogen: Pathogenesis, virulence factor
- Clinical manifestation: history + physical examination
- Diagnosis criteria: clinical +/- laboratory
- Treatment and prevention

Aerobic

Gram negative bacilli

Glucose fermenter

Oxidase

-

+

Lactose fermenter

A/A(G)

- *Escherichia*
- *Klebsiella*
- *Enterobacter*

Non-lactose fermenter

K/A

- *Salmonella*
- *Shigella*
- *Proteus*
- *Serratia*
- *Yersinia*

- *Vibrio*
- *Aeromonas*
- *Plesiomonas*

Non-Glucose fermenter

- *Pseudomonas*
- *Burkholderia*
- *Acinetobacter*
- *Stenotrophomonas*

Fastidious (cocco)bacilli

- *Haemophilus*
- *HACEK*
- *Bordetella*
- *Pasteurella*
- *Brucella*
- *Capnocytophaga*
- *Legionella*
- *Francisella*
- *Bartonella*

S-shape Microaerophile


- *Helicobacter*
- *Campylobacter*

	Morphology	Oxidase	10% Glucose	10% Lactose
<i>Pseudomonas aeruginosa</i>	Slender bacilli	+	A	K
<i>Burkholderia pseudomallei</i>	Safety pin	+	A	A
<i>Acinetobacter baumannii</i>	Coccobacilli (pair)	-	A	A
<i>Stenotrophomonas maltophilia</i>	Bacilli	-	K	K


Pseudomonas aeruginosa

- Gram negative bacilli (slender shaped)
- Green pigmented colony
- Grape-like odor
- Nosocomial > community-acquired
 - Respiratory tract infection
 - UTI
 - Sepsis
 - Otitis externa
 - Keratitis
 - Cellulitis
 - Necrotizing Fasciitis

All systems



DOI:10.3390/vaccines10071100




A)

- Examples of different pigments and pigment combinations produced in Mueller-Hinton agar with antibiotics;

B)

- 1: **pyoverdine**,
- 2: fluorescein,
- 3: fluorescein + **pyocyanin**,
- 4: pyomelanin,
- 5: pyoverdine + pyocyanin,
- 6: pyoverdine + fluorescein,
- 7: pyoverdine + pyomelanin,
- 8: no pigment.



Ecthyma gangrenosum

Clinical significance

- Crusted sores beneath which ulcers

Pathogen

- *Pseudomonas aeruginosa*
- Etc. *E. coli*, *K. pneumoniae* and *Candida spp.*

Pathogenesis

- Pseudomonas septicemia
- Invasion of the walls of arteries and veins leading to vascular necrosis (painless)

Risk factor

- immunocompromised
- Burns, ICU


Diagnosis


- Clinical diagnosis
- Blood culture



Treatment

- Ceftazidime iv
(antipseudomonal penicillins)





Malignant (necrotizing) Otitis externa

Risk factor


- Eldery
- DM



Treatment

- Ciprofloxacin
- Steroid

Pseudomonas aeruginosa



Anti-pseudomonal ATB


CAP TAP For Pseudo

	Drug/Class
C	Cefepime / Ceftazidime (Cephalosporins), Carbapenam except ertapenam
A	Aztreonam
P	Piperacillin–tazobactam
T	Ticarcillin
A	Aminoglycosides except Streptomycin
P	Polymyxins
F	Fluoroquinolones (Cipro, Levo)

Burkholderia pseudomallei

Risk factor

- Soil, water in North-East Thailand
- Skin eruption, wound
- Inhalation of dust
- Immunocompromise host :
 - DM,
 - CKD,
 - Thalassemia,





Virulence factor

- T3SS, T6SS
- LPS
- Exotoxin + Extracellular enzymes; dermatonecrotic toxin
- Hemolysin?
- Mortality enhancing polypeptide?
- Iron-acquisition : siderophore (malleobactin)

Burkholderia pseudomallei

- Suppurative Parotitis
 - No U/D, No septicemia, prior Tx as *S. aureus*



- Pneumonia
- Liver abscess



48-72hrs

- Earthy/Musty odor



Figure 221-1 Natural history of infection with *Burkholderia pseudomallei*. GIT, gastrointestinal tract.

Burkholderia pseudomallei

Laboratory diagnosis


- Culture: Pus, Hemoculture
- Serology diagnosis:
 - Indirect Hemagglutination (IHA): Antibody
 - Lateral flow assay (LFA): Antigen
- Molecular diagnosis: PCR

IHA Titer	Interpretation (in Thailand)
<1:40	Usually considered negative
≥1:160	May support diagnosis in symptomatic patients
≥1:640 or rising titer	Strongly suggestive of active infection, especially with clinical signs
Stable high titer	May reflect previous exposure, not necessarily active disease

- single high titer is not diagnostic in endemic areas. Seroconversion or a fourfold rising
- Up to 60% of healthy adults may have IHA titers ≥1:40

Treatment

- Induction (10 – 14 days)
 - Ceftazidime 2 g IV q 8 hr
- Maintenance (20 weeks)
 - Cotrimoxazole (80/400) 2 x 2 or
 - Amoxiclav (1 g) 2 x 2



IHA



Active Melioidosis Detect™ (LFA)

Vibrio vulnificus

+ Virulence factor

- Capsule
- Hemolysins and Cytolysins:
- Metalloproteases
- Siderophores

Risk factors





- After trauma and exposure to marine animals
- Consumed raw oysters within 7 days

Clinical significance

- Sudden onset of fever and chills,
- Cellulitis or NF (Hemorrhagic bleb)
- Mimic gas gangrene wound
- Septicemia - 50% death
- Diarrhea and Vomiting (less)

Treatment and prevention

- 3rd gen cephalosporin IV
- Debridement






Necrotizing Fasciitis (Hemorrhagic bleb)

Aeromonas spp.



Species

- *Aeromonas hydrophila* - wound
- *Aeromonas dhakensis* - formly hydrophila
- *Aeromonas caviae* - GI
- *Aeromonas veronii* - wound/GI- leech




Risk factor

- *Contact or consumption contaminated fresh water*

Clinical significance

- Gastroenteritis (self limiting)
- Cellulitis to necrotizing fasciitis.
- Bacteremia and Sepsis



Virulence factor

- Enterotoxins and Cytotoxins
- Hemolysins.
- Adhesins
- Type III Secretion System (T3SS)



Treatment

- Supportive
- Ciprofloxacin or Ceftriaxone
- Debridement



Necrotizing fasciitis

Clinical significance

- flesh-eating disease, rapid progression

Pathogen

- Type I (Polymicrobial) 80%; aerobic and anaerobic
- Type II; *S. pyogenes*, *S. aureus*
- Type III; *V. vulnificus*, *A. hydrophila*
- Type IV; fungus eg, *Candida spp.*

Pathogenesis

- Rapid Bacterial Proliferation
- Vascular thrombosis,
- Tissue Ischemia and Necrosis
- Systemic Toxicity; Multi-organ failure

Diagnosis

- Clinical diagnosis, Imaging, Blood culture
- Pus, Tissue Gram stain, culture



Treatment

- Surgical Debridement
- broad-spectrum intravenous antibiotics



Khon Kaen University

Thank You



Any questions?

PARABUD@KKU.AC.TH

